

Scotland's Rural College

Genetic parameters of subclinical macromineral disorders and major clinical diseases in post parturient Holstein cows

Tsiamadis, V; Banos, G; Panousis, N; Kritsepi-Konstantinou, M; Arsenos, G; Valergakis, GE

Published in:
Journal of Dairy Science

DOI:
[10.3168/jds.2015-10789](https://doi.org/10.3168/jds.2015-10789)

First published: 07/09/2016

Document Version
Peer reviewed version

[Link to publication](#)

Citation for pulished version (APA):

Tsiamadis, V., Banos, G., Panousis, N., Kritsepi-Konstantinou, M., Arsenos, G., & Valergakis, GE. (2016). Genetic parameters of subclinical macromineral disorders and major clinical diseases in post parturient Holstein cows. *Journal of Dairy Science*, 99(11), 4 - 7. <https://doi.org/10.3168/jds.2015-10789>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Genetic Parameters of Subclinical Macromineral Disorders and Major
Clinical Diseases in Post Parturient Holstein Cows**

**V. Tsiamadis, * G. Banos*§, N. Panousis, † M. Kritsepi-Konstantinou, ‡ G. Arsenos, * G.
E. Valergakis***

*Department of Animal Production, Faculty of Veterinary Medicine, Box 393, Aristotle University of
Thessaloniki, GR-54124 Thessaloniki, Greece

§Scotland's Rural College/Roslin Institute, Edinburgh, Scotland, UK EH25 9RG

†Clinic of Farm Animals, Department of Clinics, Faculty of Veterinary Medicine, Aristotle University of
Thessaloniki, GR-54124 Thessaloniki, Greece

‡Diagnostic Laboratory, Department of Clinics, Faculty of Veterinary Medicine, Aristotle University of
Thessaloniki, GR-54124 Thessaloniki, Greece

¹Corresponding author: Georgios E. Valergakis, phone: +30 2310 99 98 50, fax: 2310 99 98 92,
email: geval@vet.auth.gr

Interpretive Summary

The genetic parameters of subclinical hypocalcemia, hypophosphatemia, subclinical hypomagnesemia, hypokalemia and hyperphosphatemia during the first 8 days after calving were studied in Holstein dairy cows. Repeated measurements of calcium, phosphorus, magnesium and potassium serum concentrations together with recordings of clinical diseases during the same period were used in random regression model analyses. The heritability estimates of the associated health traits suggest that genetic selection is feasible and could help minimize health problems after calving.

ABSTRACT

The main objective of this study was to assess the genetic parameters of subclinical disorders associated with subclinical hypocalcemia (**SCHCa**), hypophosphatemia (**HypoP**), subclinical hypomagnesemia (**SCHMg**), hypokalemia (**HypoK**) and hyperphosphatemia (**HyperP**), as well as of major clinical diseases after calving in Holstein cows. The secondary objective was to estimate the associated genetic and phenotypic correlations among these subclinical and clinical conditions after calving in Holstein cows. The study was conducted in 9 dairy herds located in Northern Greece. None of the herds used any kind of preventive measures for milk fever (**MF**). A total of 1,021 Holstein cows with pedigree information were examined from November 2010 until November 2012. The distribution across parities was 466 (parity 1), 242 (parity 2), 165 (parity 3) and 148 (parity 4 and above) cows. All cows were subjected to a detailed clinical examination and blood sampled on the 1st, 2nd, 4th and 8th day after calving. Serum concentrations of Ca, P, Mg and K were measured in all samples, while β -hydroxybutyrate acid (BHBA) was measured only for day 8. The final data set included 4,064 clinical and 16,848 biochemical records (4,020 Ca, 4,019 P, 4,020 Mg and 3,792 K and 997 BHBA). Data of 1,988 observations of Body Condition Score (BCS) at days 1 and 8, were also available. All health traits were analyzed with a univariate random regression model. The genetic analysis for macromineral-related disorders included 986 cows with no obvious signs of MF (35 cows with MF were excluded). Analysis for other health traits included all 1,021 cows. A similar single record model was used for the analysis of BHBA. Genetic correlations among traits were estimated with a series of bivariate analyses. Statistically significant daily heritabilities of SCHCa (0.13 – 0.25), HypoP (0.18 – 0.33), SCHMg (0.11 – 0.38) and HyperP (0.14 – 0.22) were low to moderate, while that of HypoK was low (0.08 – 0.10). The heritability of BCS was 0.20 ± 0.10 . Statistically

significant daily heritabilities of clinical diseases were those of MF (0.07 – 0.11), left displaced abomasum (0.19 – 0.31) and mastitis (0.15 – 0.41). Results suggest that these health disorders are heritable traits and could be minimized with proper genetic selection. Statistically significant phenotypic correlations were estimated for the first time between macromineral concentrations and almost all transition cow metabolic and infectious health disorders.

Key words: subclinical macromineral disorders, postpartum diseases, genetic parameters

INTRODUCTION

During the transition period (3 weeks before to 3-4 weeks after calving) the modern high producing dairy cow is at increased risk of encountering a multitude of interrelated health disorders (Larsen et al., 2001; Lean et al., 2013). In a study that included 151,000 records, Ingvarsten et al. (2003) clearly demonstrated that disease incidence is highest during the first 10 days after calving. Negative energy balance, macromineral-related disorders and reduced immunity are the three major causes of transition period diseases (Goff, 2006a). Prevention of health disorders around calving is based on the implementation of various managerial and nutritional strategies; for example, body condition score (BCS) evaluation and post calving β -hydroxybutyric acid (BHBA) serum concentration are proposed to be routinely used as energy balance indicators (Oikonomou et al., 2008a; LeBlanc, 2010).

Macromineral serum concentration changes are mainly caused by increased cow requirements at the onset of lactation combined with reduced feed intake and possibly delayed homeostatic mechanisms (Goff, 2006a). Macromineral-related disorders, relating to calcium (Ca), phosphorus

(P), magnesium (Mg) and potassium (K) concentrations, are at the center of the disease cascade that dairy cows experience during the transition period (Goff, 2004), in either clinical or subclinical form (Goff, 2006b).

Subclinical hypocalcemia (SCHCa, serum Ca concentration < 8.3 mg/dL) is by far the most common macromineral-related health disorder associated with calving (Horst and Goff, 2003; Goff, 2008; Peek and Divers, 2008). Clinical hypocalcemia (parturient paresis – “milk fever”, MF) has a detrimental role in major post-calving clinical disease incidence, since it is associated with: retained fetal membranes (RFM), metritis (MET), mastitis (MAST), displaced abomasum (left or right, LDA and RDA, respectively), ketosis (KET) and uterine prolapse (UP) (Correa et al., 1990; Gröhn and Bruss, 1990; DeGaris and Lean, 2008). Subclinical hypocalcemia is assumed to have the same negative effects but relevant literature is lacking.

Lower than normal P concentrations (HypoP, P < 4.2 mg/dL) are common at the onset of lactation; recumbent MF dairy cows often have very low P concentration (P < 2.0 mg/dL) (Goff, 2004). Elevated P concentrations (HyperP, P > 7.80 mg/dL) increase the risk of MF (Lean et al., 2013; Grünberg, 2014). While clinical hypomagnesaemia (“grass tetany”, serum Mg < 1.0 mg/dL) may still appear in grazing herds, it is not at all common in confined and TMR-fed cows (Peek and Divers, 2008). On the other hand, subclinical hypomagnesemia (SCHMg, serum Mg < 1.8 mg/dL) is involved in the etiology of SCHCa and MF (Littledike et al., 1983; Rude, 1998; Schonewille et al., 2008). Mild hypokalemia (serum K between 2.6 and 3.8 mmol/L) is common in early lactation (Sattler and Fecteau, 2014), while severe hypokalemia (serum K < 2.5 mmol/L)

is very rare in dairy cattle, mostly associated with concurrent infectious disease (Sattler et al. 1998).

Macromineral-related disorders usually resolve by the end of the first week post-calving but their effects are long-lasting, impairing milk production and reproductive efficiency of dairy cows (Goff, 2006b). Despite the extensive knowledge regarding the pathophysiology of macromineral-related disorders and the various management practices that may alleviate them (Thilising-Hansen et al., 2002; Goff, 2004; Mulligan et al., 2006), problems are still common. Disease incidence rates, even in many well-managed herds, still remain unacceptably high (Mulligan and Doherty, 2008). During the last decades, genetic selection for disease resistance enjoys increased popularity because genetic progress, no matter how small, is permanent and cumulative (Eggen, 2012). Genetic parameters for various clinical diseases around calving have been estimated in several large scale studies (Lin et al., 1989; Lyons et al., 1991; Heringstad et al., 2005). Heritabilities of Ca, P, Mg and K serum concentrations have only recently been reported (Tsiamadis et al., 2016); however, there is lack of information concerning subclinical macromineral-related disorders.

The objectives of this study were to estimate: 1) the heritability of SCHCa, HypoP, HyperP, SCHMg, HypoK, BHBA and BCS, 2) the heritability of major clinical health disorders (MF, RFM, MET, MAST, LDA, RDA, KET and UP) and 3) relevant genetic and phenotypic correlations, during the first 8 days after calving.

MATERIALS AND METHODS

The research was conducted in compliance with institutional guidelines and approved by the Research Committee of the Aristotle University of Thessaloniki, Thessaloniki, Greece. All farmers gave informed consent for the cows to be included in the study and to undergo the testing procedures.

Animals and Management

A total of 1,021 Holstein cows from 9 commercial free-stall dairy herds in Northern Greece were included in the study. The distribution across parities was 466, 242, 165 and 148 cows for parities 1, 2, 3 and 4 and above, respectively. Farms were visited regularly between November 2010 and November 2012 for data collection. No herd used any kind of preventive measures for hypocalcemia. Total mixed rations (TMR) were formulated to meet or exceed net energy and metabolizable protein requirements according to National Research Council recommendations (NRC, 2001).

Clinical Examination, Blood Sampling and Analyses

All animals were clinically examined and blood sampled by the first author on the 1st, 2nd, 4th and 8th day after calving. Body condition score was recorded on the 1st and 8th day after calving using the 1- to 5- point scale of Ferguson et al. (1994), in increments of 0.25. At this scale, 1 is for emaciated and 5 for obese animals.

Blood sampling was performed by coccygeal venipuncture into 10-ml vacuum glass tubes without anticoagulant (BD Vacutainer®, Plymouth, United Kingdom) for serum macromineral

measurements. Samples were placed in a cooler, transported to the Diagnostic Laboratory of the Faculty of Veterinary Medicine and centrifuged immediately upon arrival (3,000 x g for 15 min, room temperature 21°C). Serum was transferred into polyethylene tubes and stored at -80°C until assay. All sera were analyzed for total Ca and Mg concentrations using flame atomic absorption spectrophotometry (Perkin ElmerAAAnalyst 100, Perkin Elmer Co, Norwalk, CT, USA), according to manufacturer's instructions. Serum inorganic phosphorus concentrations were determined photometrically using a Flexor E autoanalyzer (Vital Scientific, Spankeren, The Netherlands), according to the procedure described by Daly and Ertingshausen (1972), with the use of standard commercial reagents (Thermo Fisher Scientific Inc. USA). Potassium serum concentrations were measured using an ion-selective electrode according to manufacturer's instructions (Electrolyte Analyzer 9180, Roche Austria). The intra- and inter-assay coefficients of variation for all the above analyses were less than 3%. Beta-hydroxybutyric acid was measured only on the 8th day after calving by a spectrophotometric kinetic method (Bruss, 2008). The intra-assay coefficient was 2 to 4%, while the inter-assay coefficient was 4 to 8%, both of which are within the desirable range.

Disease Definitions and Cut-offs

In our study, SCHCa, HypoP, HyperP, SCHMg and HypoK were defined based on threshold values provided in relevant literature and were expressed as presence or absence of the condition (binary traits). Animals with serum concentrations below or equal to 8.3 mg/dL for Ca, 4.2 mg/dL for P, 1.8 mg/dL for Mg, and 3.9 mmol/L for K, were considered as cases of SCHCa, HypoP, SCHMg and HypoK, respectively (Goff, 2008; Divers and Peek, 2008; Horst and Goff, 2003). Moreover, animals with inorganic serum P concentration ≥ 7.80 mg/dL were considered

HyperP cases, while cows with serum BHBA $\geq 1,200$ $\mu\text{mol/L}$ were considered subclinically ketotic (Divers and Peek, 2008).

Clinical diseases were defined as follows: a) MF, standing (showing mild ataxia, excitability, muscle tremors and reduced ruminal motility) or recumbent cow (Kelton et al., 1998; Oetzel, 2011); b) RFM, fetal membranes were visible at the vulva or were identified in the uterus by vaginal examination more than 12 hours after calving (Melendez et al., 2003); c) MET, fetid uterine discharge, with or without fever (Sheldon et al., 2006); d) MAST, milk clots or abnormal mammary discharge from one or more quarters (Kelton et al., 1998); e) KET, decreased appetite together with elevated blood BHBA ($> 2,000$ $\mu\text{mol/L}$), in the absence of obvious concurrent disease (Kelton et al., 1998; Duffield et al., 2009); f) LDA/RDA, decreased appetite accompanied by a clearly audible “ping” sound, produced by percussion of the left/right abdominal wall (between the 9th and 12th ribs), respectively (Kelton et al., 1998).

Data set

Pedigree information was available for all 1,021 cows (332 common sires and 786 common dams). The total population in the study increased to 4,262 animals, when all available pedigree information included, spanning the last 5 (overlapping) generations. Calving date, parity number, calving ease and twinning was recorded. From the 1,021 cows, 35 were diagnosed with MF during the first 4 days after calving, treated appropriately with intravenous Ca and excluded from the genetic analysis of macromineral-related health traits. Therefore, 986 cows were included in the genetic analysis for SCHCa, HypoP, HyperP, SCHMg and HypoK. However, genetic

analysis for the other recorded clinical health traits (MF, RFM, MET, MAST, LDA, RDA, KET and UP) included all 1,021 cows.

The final data set included 4,064 clinical observations for MF, RFM, MET, MAST, LDA, RDA, KET and UP. Moreover, observations for death (DE) and involuntary culling (INVCULL) during the same time-period were also included in the data set, as well as 1,988 BCS records. In total, 16,848 biochemical records were available, consisting of 4,020 Ca, 4,019 P, 4,020 Mg, 3,792 K (days 1, 2, 4 and 8 after calving) and 997 BHBA (only on day 8) measurements. Changes of the macrominerals concentrations between day 1 and day 4, as well as between day 1 and day 8 were calculated as the regression slope of macromineral concentrations on time. Thus, these measurements reflected the average daily change in said concentrations and were treated as different traits.

Statistical Analysis

Macromineral-related and disease-related health traits measured on days 1 through 8 were analyzed with a random regression model which accounted for the covariance between successive records of the same animal; each trait was analyzed separately:

$$Y_{ijkmn} = HYS_i + L_j + M_k + a_1 \cdot age + \sum_{m=0}^2 b_m P_m D + \sum_{m=0}^2 A_{nm} P_m D + e_{ijkmn}$$

(1)

where:

Y_{ijkmn} is the health trait record of cow n ;

HYS_i is the fixed effect of herd-year-season of calving i (72 levels);

205 L_j the fixed effect of number of lactation j (4 levels);
 206 M_k the fixed effect of calendar month k (12 levels);
 207 a_1 the linear regression coefficient on age at calving (age);
 208 P_m orthogonal polynomial of order m ;
 209 b_m the fixed regression coefficient on days from calving (D);
 210 A_{nm} the random regression coefficient on days from calving associated with the additive
 211 genetic effect of cow n including all pedigree data (4,262 animals spanning five
 212 generations);
 213 e_{ijkmn} the random residual term.

214 The fixed effects in the model including the polynomial order in the fixed regression were fitted
 215 after preliminary analyses had confirmed their statistically significant effect ($P < 0.05$) on the
 216 traits based on the F-test. Further increasing the order of the polynomial did not have a
 217 significant effect ($P > 0.05$). Similarly, the final order of the random polynomial (third for either
 218 trait) was determined with the use of the log-likelihood ratio test in sequential analyses of
 219 gradually increasing orders. The final order choice was also confirmed with the Akaike
 220 Information Criterion test. Four measurement error classes were defined for each the day from
 221 calving (1st, 2nd, 4th and 8th). The definition of these classes, even at this small time span, aimed
 222 to capture the day-to-day differences in health events at the beginning of lactation. Covariances
 223 between the error classes were assumed to be zero.

224

225 A random permanent environment effect was also fitted to model (1) resulting in a practically
 226 zero corresponding variance component estimate, possibly due to the short period our data

spanned (8 days). The log-likelihood ratio tests between the models including and excluding permanent environment were not significant ($P>0.05$) in all analyses.

There was also an effort to fit a Logit function in model (1) to account for the binary nature of the disease traits. However, this was proved unfeasible within the context of a random regression model.

Serum BHBA concentration for day 8 from calving and average estimates for BCS on days 1 and 8 and serum concentration changes between day 1 and day 4 (days 1-4), as well as day 1 and day 8 (days 1-8) after calving were analyzed using the following model:

$$Y_{ijkm} = HYS_i + L_j + a_1 \cdot age + A_k + e_{ijkm} \quad (2)$$

where Y_{ijkm} is the log-transformed value for serum BHBA concentration or BCS or macromineral concentration change of cow k; A_k is the additive genetic effect of cow k and all effects are as in model 1.

Estimates of variance components from each model were used to calculate heritabilities for each trait, with the following equation:

$$h^2 = \frac{\sigma_a^2}{\sigma_p^2}$$

where h^2 = the heritability estimate, σ_a^2 = the additive genetic variance and σ_p^2 = the phenotypic variance.

Genetic (r_a) and phenotypic (r_p) correlations among all traits analyzed with the above models were estimated based on co-variance components derived with a series of bivariate analyses based on the same model described for each trait, with the following equation:

$$r_{(a,p)} = \frac{Cov_{(a,p)}(X, Y)}{\sqrt{\sigma_{(a,p)X}^2 \times \sigma_{(a,p)Y}^2}}$$

where $Cov_{(a,p)}(X, Y)$ = the additive genetic (Cov_a) or phenotypic (Cov_p) co-variance of traits X and Y and $\sigma_{(a,p)X}^2$ and $\sigma_{(a,p)Y}^2$ are the genetic (σ_a^2) or phenotypic (σ_p^2) variances of relevant traits.

All analyses were conducted using the statistical software package ASREML (Gilmour and Gogel, 2006). In all cases, statistical significance was set at $P < 0.05$.

RESULTS

Descriptive statistics for Ca, P, Mg, K and BHBA serum concentrations and BCS are presented in Table 1. Average incidence of health disorders during the same time period after calving is presented in Table 2.

Random regression model was used for the generation of prevalence curves across all lactations for all health disorders during the first 8 days after calving. However, this was possible only for

SCHCa, HypoP, SCHMg, HypoK, and HyperP (Figure 1), and also for MF, LDA and MAST (Figure 2). The remaining health disorders had either low incidence (RDA, UP, INVCULL, DE), or were not present throughout the entire 8 day period (RFM: present only the first day; MET, KET: present mainly after the 4th day), thus rendering it impossible to generate curves.

Day-to-day variances (phenotypic, genetic, and residual) and heritabilities for SCHCa, HypoP, SCHMg, HypoK, and HyperP are shown in Table 3 and for MF, LDA, and MAST in Table 4. All estimates presented were statistically greater than zero ($P < 0.05$). Day-to-day heritability estimates were low to moderate for SCHCa ($h^2 = 0.13 - 0.25$), HypoP ($h^2 = 0.18 - 0.33$), HyperP ($h^2 = 0.14 - 0.22$), SCHMg ($h^2 = 0.11 - 0.38$) and LDA ($h^2 = 0.19 - 0.31$), low for HypoK and MF ($h^2 = 0.07 - 0.11$), and moderate to high for MAST ($h^2 = 0.15 - 0.41$). Regarding serum BHBA, the heritability estimate was not statistically significant ($h^2 = 0.073 \pm 0.077$, $P = 0.12$), while for BCS was statistically significant ($h^2 = 0.20 \pm 0.10$, $P < 0.05$).

Significant genetic correlations: a) between serum Ca, P, Mg and K concentrations and health disorders, b) of macromineral concentration changes in days 1-4 and 1-8 after calving with health disorders, and c) among health disorders were not detected in the present study.

Statistically significant phenotypic correlations between overall serum Ca, P, Mg and K concentrations and health disorders during the first 8 days after calving are shown in Table 5. calcium, Mg and K concentrations had high negative correlations with the related subclinical disorders; this was not the case with P. Serum Ca concentrations had a low positive correlation with BCS and a low negative correlation with BHBA; moreover, correlations with most health

disorders were negative, either low (HypoP, HypoK, HyperP, LDA, RFM, MET and DE) or moderate (MF, SCHMg). Correlations of Mg and K concentrations with health disorders were similar with those of Ca. Magnesium (but not K) concentrations had a low positive correlation with BCS. For those health disorders that significant correlations were detected, all were negative albeit low. Regarding P, only a high positive correlation with HyperP and low ones with MAST and UP were detected.

Statistically significant phenotypic correlations of serum macromineral concentrations on day 1 and their changes from day 1 to 4 and 1 to 8 after calving with health disorders during the first 8 days after calving are shown in Table 6. Calcium concentrations on day 1 and their changes had similar correlations with the various health disorders as those presented in Table 5. Calcium concentration on day 1 was mostly correlated with low concentrations of the other macrominerals, with Ca-related disorders (SCHCa and MF) and MET, while Ca changes were correlated with RFM, MET, KET,DE and INVCULL. Phosphorus concentration on day 1 had similar correlations with the same health disorders as those presented in Table 5, as well. Moreover, a negative correlation with BCS was detected. Phosphorus decrease over time was negatively correlated with HyperP and positively correlated with MF, RFM and DE. High Mg concentration on day 1 was again positively correlated with BCS and negatively with SCHMg and MET. Magnesium changes were correlated with SCHCa, HypoP and BHBA, LDA, MET and DE. Potassium concentrations on day 1 had also similar correlations with the same health disorders as those presented in Table 5. Potassium changes were significantly correlated with HyperP and SCHCa.

Statistically significant phenotypic correlations of MF, SCHCa, HypoP, HyperP, SCHMg and HypoK with transition period health events are shown in Table 7. Correlations were low but follow the same pattern as those of the respective macromineral serum concentrations, definitely connecting these health conditions with each other.

DISCUSSION

This study aimed to estimate genetic parameters of subclinical and clinical diseases that occur during the first 8 days after calving. Detailed records were obtained including day-to-day clinical examination of cows by the same veterinarian.

Incidence of health disorders was estimated during the first 8 days after calving and a mixed model was used for the estimation of the day-to-day prevalence, which was modeled as a third polynomial fixed regression on days postpartum. The latter gave an accurate mapping of the health status of the population in study.

Prevalence in this study is in agreement with those of Reinhardt et al. (2011) regarding SCHCa, of Staufenbiel (2002) and Macrae et al. (2006) regarding HypoP and HyperP, of Masoero et al. (2003) regarding SCHMg and of Peek and Divers (2008) regarding HypoK. Moreover, incidence and prevalence of major clinical diseases recorded in this study were very similar with those reported in the literature (Kelton et al., 1998; Heringstad et al., 2005; Melendez and Risco, 2005; LeBlanc, 2008). Therefore, our estimations of various genetic parameters are concurrent with the global Holstein population kept under similar management practices.

Heritabilities of Ca, P, Mg and K serum concentrations have only recently been reported (Tsiamadis et al., 2016). Heritabilities of SCHCa, HypoP, HyperP, SCHMg and HypoK estimated in this study, are reported for the first time in the literature. They were low to moderate but generally within the range reported for other traits such as milk yield ($h^2 = 0.20 - 0.50$ (Castillo-Juarez et al., 2000; Windig et al., 2006; Bastin et al., 2011)), somatic cell count ($h^2 = 0.03 - 0.11$ (Koeck et al., 2012; Heringstad et al., 2006)) and longevity ($h^2 = 0.01 - 0.36$ (Veerkamp and Brotherstone, 2001; Jamrozik et al., 2008)), which are already used in breeding programs. Heringstad et al. (2007) reported that there is potential for selection against metabolic disease resistance and there are several studies that investigate the genetic basis of non-infectious disease resistance (Lin et al., 1989; Lyons et al., 1991; Abdel-Azim et al., 2005). Substantial and statistically significant genetic variance estimates derived in the present study corroborate these assertions.

At the same time, low heritability estimates suggest that environmental factors have a strong influence in the etiology of the studied traits. Nutrition, management and housing of cows during the transition period emerge as critical factors for prevention of these health disorders in the short term. Nevertheless, genetic selection for resistance for these macromineral deficiency traits could be effective and add permanent benefits to successfully address the problem in the long term, thereby complementing management practices.

Heritability of BHBA in the present study was not statistically significant ($h^2 = 0.073 \pm 0.077$). Oikonomou et al. (2008b) also reported heritability estimates in primiparous Holstein cows ($h^2 = 0.25 \pm 0.18$), which were not statistically significant. However, recently, van der Drift et al.

(2012) in a study of 1,772 Holstein cows of various parities between 5 and 60 days after calving from 123 herds, using a similar animal model, reported a heritability estimate of 0.17 ± 0.06 ($P < 0.001$). This higher heritability estimate can be attributed to the much wider sampling period (1 blood sample between 5 to 60 days after calving), which possibly resulted in a higher incidence of hyperketonemia. The heritability estimate of BCS was statistically significant in the present study ($h^2 = 0.20 \pm 0.10$). Koenen et al. (2001), Veerkamp and Brotherstone (2001) and Oikonomou et al. (2008b) reported higher estimates (0.28 – 0.50) that were statistically significant. Others (Jones et al., 1999; Dechow et al., 2001; Bastin et al., 2010) have reported lower estimates (0.07 – 0.20), which are similar to our results. Heritability estimates of BCS tend to be larger in mid to late lactation (Dechow et al., 2001) and it is likely that the focus of this study on the first week after calving could have led to this moderate estimate.

The present study's estimates of MF heritability ($h^2 = 0.07 - 0.11$) are in agreement with those of Dyrendahl et al. (1972), Uribe et al. (1995), Pryce et al. (1997), Van Dorp et al. (1998) and Heringstad et al. (2005). These, however, are generally lower than estimates reported by Lin et al. (1989), Lyons et al. (1991) and Abdel-Azim et al. (2005) ($h^2 = 0.30 - 0.40$). Differences in estimates can be attributed to methodology of statistical analysis, data collection (farm records), and type and age of the population studied.

Our heritability estimates for LDA ($h^2 = 0.18 - 0.31$) are similar to those reported by Uribe et al., (1995) ($h^2 = 0.304 \pm 0.005$, across lactation with a threshold model). This is higher than other estimates from linear models reported by Lyons et al. (1991), Appuhamy et al. (2009) and Koeck et al. (2013). Moreover, Wolf (2001) and Hamann et al. (2004) with the use of threshold models

reported heritability estimates above 0.50. The moderate to high heritability estimates of the present study can be attributed to a more accurate recording of the displacement made by the veterinarian and to the binary nature of the trait that posed no ambiguity to the severity of the disease and thus to the certainty of the diagnosis.

Heritability estimates for MAST vary across studies. Lin et al. (1989) reported heritabilities of 0.19±0.08, 0.31±0.10 and 0.18±0.09 for the 1st, 2nd and 3rd+ lactation, respectively. Uribe et al. (1995) reported similar estimates for 1st lactation cows ($h^2 = 0.15 \pm 0.05$) but for all lactations estimates were zero. Zwald et al. (2004) and Heringstad et al. (2005) reported much lower estimates ($h^2 = 0.09 \pm 0.01$); more recently, Pérez-Cabal et al. (2009) and Vazquez et al. (2009) also reported similar heritabilities ($h^2 = 0.09$ and $h^2 = 0.13$, respectively), while Koeck et al. (2013) estimated the heritability of clinical mastitis at 0.02±0.004. However, all these studies estimated mastitis heritability across lactation. Our estimates ($h^2 = 0.15 - 0.41$) cover a small portion of the entire lactation, only the first 8 days. Considering that clinical mastitis immediately after calving is influenced by factors such as dry period management and compromised immune status due to calving (Kimura et al., 2006; LeBlanc, 2010), this may well be a different trait which, based on our results, could potentially respond to selection.

The present study did not detect any significant genetic correlation of Ca, P, Mg and K serum concentrations and BCS with any postpartum health disorders. The absence of genetic correlations could be attributed to the multifactorial etiology of most of these health events: infectious agents may co-exist with metabolic and managerial deficiencies. Moreover, this lack of genetic correlation may support the idea that these traits are controlled genetically by different

genes and individual selection should be applied. Contrary to expectations, this study did not find a significant genetic correlation between SCHCa and MF. However, considering the disease definitions, MF cases were defined as standing (showing mild ataxia, excitability, muscle tremors and reduced ruminal motility) or recumbent cows; therefore, MF definition was solely based on symptoms and not in any serum Ca measurement. Furthermore, this absence of genetic correlation could also be attributed to the low incidence of MF. In this study, recumbent cows were immediately treated with intravenous Ca solutions, rendering the measurement of serum Ca concentrations meaningless. Moreover, it is known that there is no specific threshold of Ca serum concentrations which always results in recumbent cows. Regarding the absence of any genetic correlation of the remaining macrominerals with other health disorders, this may also be attributed to the multifactorial etiology and to the low incidence of some of the health disorders (e.g. metritis, mastitis and ketosis). On the other hand, the lack of any significant genetic correlation in this study may be incidental. Therefore, as this is the first study of its kind, the genetic analysis of other independent data sets may shed more light on this issue; more research is needed in order to clarify these issues.

The reported phenotypic association of clinical and subclinical hypocalcemia with various diseases after calving is based almost solely on pathophysiology, because of calcium's central metabolic role; it is generally assumed that P and Mg serum concentrations are associated with the same postpartum diseases through their relation with Ca metabolism (Rude, 1998; Goff, 2000; DeGaris and Lean, 2008). In a study of 2,190 cows from 33 herds, Curtis et al. (1983) showed that cows with clinical hypocalcemia (MF) were at greater risk of developing dystocia (6.5 times), RFM (3.2 times), KET (8.9 times) and MAST (8.1 times). Martinez et al. (2012)

found that cows with low serum Ca have higher BHBA concentrations. However, large scale research-based evidence for any association of subclinical macromineral-related disorders with postpartum cow health is lacking. In the present study, statistically significant phenotypic correlations of the four major macrominerals' serum concentrations and the corresponding subclinical disorders with the early postpartum disease cascade in dairy cows are reported for the first time. A strong association with energy metabolism is evident both at the KET and BHBA, as well as the BCS levels, with serious indirect and direct implications for future reproductive performance (RFM, MET and UP), MAST and replacement rates (LDA, INVCULL and DE). The correlation of HyperP with MAST is a novel finding and the exact mechanism of this association has to be further investigated. These results highlight not only the need for genetic selection against these subclinical disorders, which is feasible based on our heritability estimates, but also for enhanced implementation of pertinent management practices.

Herd management during early postpartum is a challenge for modern dairy farms. The ability of an animal to maintain normal serum macromineral concentrations is consistent with the successful management of the numerous health events after calving. Rapid metabolic changes of animals combined with stressors such as nutritional and grouping changes further compromise immunity status, favor metabolic and infectious diseases, and downgrade productivity and welfare. Postpartum health monitoring programs are implemented in many dairy farms worldwide since they greatly contribute to the early recognition and proper treatment of sick animals (Risco, 2011). Obviously, genetic selection can provide a valuable tool, as well. Standardized health monitoring programs across regions and countries could provide accurate phenotype information for novel functional traits, the discovery of their genetic markers and

finally, the creation of a new index (“disease resistance early postpartum”). This is, indeed, a very exciting prospect.

CONCLUSIONS

More research is needed on this issue, but results of the present study clearly indicate that subclinical Ca, P, Mg and K disorders during the first week after calving are heritable traits. Moreover, significant heritability estimates of BCS and MF, MAST and LDA during the same period were also derived. These genetic parameters can potentially be used to develop health indices for the selection of dairy cows that will effectively resist health challenges immediately after calving. Phenotypic correlations of high prevalence subclinical macromineral disorders with clinical diseases, reveal a deeper interrelationship among these traits and stresses the need for both innovative genetic selection and effective management practices.

REFERENCES

- Abdel-Azim, G.A., A.E. Freeman, M.E. Kehrli, S.C. Kelm, J.L. Burton, A.L. Kuck, and S. Schnell. 2005. Genetic basis and risk factors for infectious and noninfectious diseases in US Holsteins. I. Estimation of genetic parameters for single diseases and general health. *J. Dairy Sci.* 88:1199–1207. doi:10.3168/jds.S0022-0302(05)72786-7.
- Bastin, C., N. Gengler, and H. Soyeurt. 2011. Phenotypic and genetic variability of production traits and milk fatty acid contents across days in milk for Walloon Holstein first-parity cows. *J. Dairy Sci.* 94:4152–4163. doi:10.3168/jds.2010-4108.
- Bastin, C., S. Loker, N. Gengler, A. Sewalem, and F. Miglior. 2010. Genetic relationships between body condition score and reproduction traits in Canadian Holstein and Ayrshire

476 first-parity cows. *J. Dairy Sci.* 93:2215–28. doi:10.3168/jds.2009-2720.

477 Bruss, M.L. 2008. Lipids and Ketones. *In* Clinical Biochemistry of Domestic Animals. 81–115.

478 Castillo-Juarez, H., P.A. Oltenacu, R.W. Blake, C.E. McCulloch, and E.G. Cienfuegos-Rivas.

479 2000. Effect of herd environment on the genetic and phenotypic relationships among milk

480 yield, conception rate, and somatic cell score in Holstein cattle. *J. Dairy Sci.* 83:807–814.

481 doi:10.3168/jds.S0022-0302(00)74943-5.

482 Correa, M.T., C.R. Curtis, H.N. Erb, J.M. Scarlett, and R.D. Smith. 1990. An ecological analysis

483 of risk factors for postpartum disorders of Holstein-Friesian cows from thirty-two New

484 York farms. *J. Dairy Sci.* 73:1515–24. doi:10.3168/jds.S0022-0302(90)78819-4.

485 Curtis, C.R., H.N. Erb, C.J. Sniffen, R.D. Smith, P.A. Powers, M.C. Smith, M.E. White, R.B.

486 Hillman, and E.J. Pearson. 1983. Association of parturient hypocalcemia with eight

487 periparturient disorders in Holstein cows. *J. Am. Vet. Med. Assoc.* 183:559–61.

488 Daly, J.A., and G. Ertingshausen. 1972. Direct method for determining inorganic phosphate in

489 serum with the “CentrifChem”. *Clin. Chem.* 18:263–5.

490 Dechow, C.D., G.W. Rogers, and J.S. Clay. 2001. Heritabilities and Correlations Among Body

491 Condition Scores, Production Traits, and Reproductive Performance. *J. Dairy Sci.* 84:266–

492 275. doi:10.3168/jds.S0022-0302(01)74476-1.

493 DeGaris, P.J., and I.J. Lean. 2008. Milk fever in dairy cows: A review of pathophysiology and

494 control principles. *Vet. J.* 176:58–69. doi:10.1016/j.tvjl.2007.12.029.

495 Divers, T.J., and S.F. Peek. 2008. Chapter 1 - The Clinical Examination. *In* Rebhun’s Diseases of

496 Dairy Cattle (Second Edition). T.J.D.F. Peek, editor. W.B. Saunders, Saint Louis. 3–15.

497 Van Dorp, T.E., J.C. Dekkers, S.W. Martin, and J.P. Noordhuizen. 1998. Genetic parameters of

498 health disorders, and relationships with 305-day milk yield and conformation traits of

499 registered Holstein cows. *J. Dairy Sci.* 81:2264–2270. doi:10.3168/jds.S0022-
 500 0302(98)75806-0.

501 van der Drift, S.G.A., K.J.E. van Hulzen, T.G. Teweldemedhn, R. Jorritsma, M. Nielen, and
 502 H.C.M. Heuven. 2012. Genetic and nongenetic variation in plasma and milk β -
 503 hydroxybutyrate and milk acetone concentrations of early-lactation dairy cows. *J. Dairy Sci.*
 504 95:6781–6787. doi:10.3168/jds.2012-5640.

505 Duffield, T.F., K.D. Lissemore, B.W. McBride, and K.E. Leslie. 2009. Impact of
 506 hyperketonemia in early lactation dairy cows on health and production. *J. Dairy Sci.*
 507 92:571–580. doi:10.3168/jds.2008-1507.

508 Dyrendahl, I., B. Henricson, and G. Jönsson. 1972. Clinical puerperal paresis and hypocalcaemia
 509 in cattle. A statistical and genetic investigation. *Zentralbl. Veterinarmed. A.* 19:621–638.

510 Eggen, A. 2012. The development and application of genomic selection as a new breeding
 511 paradigm. *Anim. Front.* 2:10–15. doi:10.2527/af.2011-0027.

512 Gilmour, A.R., Gogel, B.J., Cullis, B.R., Thompson, R. ASReml User Guide Release 2.0. VSN
 513 International Ltd., Hemel Hempstead, UK; 2006. <http://www.vsnl.co.uk/>.

514 Goff, J.P. 2000. Pathophysiology of calcium and phosphorus disorders. *Vet. Clin. North Am.*
 515 *Food Anim. Pract.* 16:319–337, vii.

516 Goff, J.P. 2004. Macromineral disorders of the transition cow. *Vet. Clin. North Am. - Food Anim.*
 517 *Pract.* 20:471–494. doi:10.1016/j.cvfa.2004.06.003.

518 Goff, J.P. 2006a. Major advances in our understanding of nutritional influences on bovine health.
 519 *J. Dairy Sci.* 89:1292–1301. doi:10.3168/jds.S0022-0302(06)72197-X.

520 Goff, J.P. 2006b. Macromineral physiology and application to the feeding of the dairy cow for
 521 prevention of milk fever and other periparturient mineral disorders. *Anim. Feed Sci.*

522 *Technol.* 126:237–257. doi:10.1016/j.anifeedsci.2005.08.005.

523 Goff, J.P. 2008. The monitoring, prevention, and treatment of milk fever and subclinical
524 hypocalcemia in dairy cows. *Vet. J.* 176:50–57. doi:10.1016/j.tvjl.2007.12.020.

525 Gröhn, Y.T., and M.L. Bruss. 1990. Effect of diseases, production, and season on traumatic
526 reticuloperitonitis and ruminal acidosis in dairy cattle. *J. Dairy Sci.* 73:2355–2363.
527 doi:10.3168/jds.S0022-0302(90)78918-7.

528 Grünberg, W. 2014. Treatment of Phosphorus Balance Disorders. *Vet. Clin. North Am. - Food*
529 *Anim. Pract.* 30:383–408. doi:10.1016/j.cvfa.2014.03.002.

530 Hamann, H., V. Wolf, H. Scholz, and O. Distl. 2004. Relationships between lactational incidence
531 of displaced abomasum and milk production traits in German Holstein cows. *J. Vet. Med.*
532 *Ser. A Physiol. Pathol. Clin. Med.* 51:203–208. doi:10.1111/j.1439-0442.2004.00626.x.

533 Heringstad, B., Y.M. Chang, D. Gianola, and G. Klemetsdal. 2005. Genetic analysis of clinical
534 mastitis, milk fever, ketosis, and retained placenta in three lactations of Norwegian red
535 cows. *J. Dairy Sci.* 88:3273–3281. doi:10.3168/jds.S0022-0302(05)73010-1.

536 Heringstad, B., D. Gianola, Y.M. Chang, J. Odegård, and G. Klemetsdal. 2006. Genetic
537 associations between clinical mastitis and somatic cell score in early first-lactation cows. *J.*
538 *Dairy Sci.* 89:2236–44. doi:10.3168/jds.S0022-0302(06)72295-0.

539 Heringstad, B., G. Klemetsdal, and T. Steine. 2007. Selection responses for disease resistance in
540 two selection experiments with Norwegian red cows. *J. Dairy Sci.* 90:2419–2426.
541 doi:10.3168/jds.2006-805.

542 Horst, R., and J. Goff. 2003. Prevalence of Subclinical Hypocalcemia in US Dairy Operations. *J.*
543 *Dairy Sci.* 86 (Suppl.:1, 247.

544 Ingvarsen, K.L., R.J. Dewhurst, and N.C. Friggens. 2003. On the relationship between

545 lactational performance and health: Is it yield or metabolic imbalance that cause production
546 diseases in dairy cattle? A position paper. *Livest. Prod. Sci.* 83:277–308.
547 doi:10.1016/S0301-6226(03)00110-6.

548 Jamrozik, J., J. Fatehi, and L.R. Schaeffer. 2008. Comparison of models for genetic evaluation of
549 survival traits in dairy cattle: a simulation study. *J. Anim. Breed. Genet.* 125:75–83.
550 doi:10.1111/j.1439-0388.2007.00712.x.

551 Jones, H., I. White, and S. Brotherstone. 1999. Genetic evaluation of Holstein Friesian sires for
552 daughter condition-score changes using a random regression model. *Anim. Sci.* 68.

553 Kelton, D.F., K.D. Lissemore, and R.E. Martin. 1998. Recommendations for recording and
554 calculating the incidence of selected clinical diseases of dairy cattle. *J. Dairy Sci.* 81:2502–
555 2509. doi:10.3168/jds.S0022-0302(98)70142-0.

556 Kimura, K., T. a Reinhardt, and J.P. Goff. 2006. Parturition and hypocalcemia blunts calcium
557 signals in immune cells of dairy cattle. *J. Dairy Sci.* 89:2588–2595. doi:10.3168/jds.S0022-
558 0302(06)72335-9.

559 Koeck, A., F. Miglior, J. Jamrozik, D.F. Kelton, and F.S. Schenkel. 2013. Genetic associations of
560 ketosis and displaced abomasum with milk production traits in early first lactation of
561 Canadian Holsteins. *J. Dairy Sci.* 96:4688–96. doi:10.3168/jds.2012-6408.

562 Koeck, A., F. Miglior, D.F. Kelton, and F.S. Schenkel. 2012. Alternative somatic cell count traits
563 to improve mastitis resistance in Canadian Holsteins. *J. Dairy Sci.* 95:432–439.
564 doi:10.3168/jds.2011-4731.

565 Koenen, E.P.C., R.F. Veerkamp, P. Dobbelaar, and G. De Jong. 2001. Genetic Analysis of Body
566 Condition Score of Lactating Dutch Holstein and Red-and-White Heifers. *J. Dairy Sci.*
567 84:1265–1270. doi:10.3168/jds.S0022-0302(01)74588-2.

568 Larsen, T., G. Møller, and R. Bellio. 2001. Evaluation of clinical and clinical chemical
 569 parameters in periparturient cows. *J. Dairy Sci.* 84:1749–1758. doi:10.3168/jds.S0022-
 570 0302(01)74610-3.

571 Lean, I.J., R. Van Saun, and P.J. Degaris. 2013. Energy and protein nutrition management of
 572 transition dairy cows. *Vet. Clin. North Am. Food Anim. Pract.* 29:337–66.
 573 doi:10.1016/j.cvfa.2013.03.005.

574 LeBlanc, S. 2010. Monitoring metabolic health of dairy cattle in the transition period. *J. Reprod.*
 575 *Dev.* 56 Suppl:S29–S35. doi:10.1262/jrd.1056S29.

576 LeBlanc, S.J. 2008. Postpartum uterine disease and dairy herd reproductive performance: A
 577 review. *Vet. J.* 176:102–114. doi:10.1016/j.tvjl.2007.12.019.

578 Lin, H.K., P.A. Oltenacu, L.D. Van Vleck, H.N. Erb, and R.D. Smith. 1989. Heritabilities of and
 579 genetic correlations among six health problems in Holstein cows. *J. Dairy Sci.* 72:180–186.
 580 doi:10.3168/jds.S0022-0302(89)79095-0.

581 Littledike, E.T., J.A. Stuedemann, S.R. Wilkinson, and R.L. Horst. 1983. Grass tetany syndrome.
 582 *In* Proceedings of John Lee Pratt International Symposium on the Role of Magnesium in
 583 Animal Nutrition. Virginia Polytechnic Institute and State University, Blacksburg, Virginia,
 584 VA, USA. 173.

585 Lyons, D.T., A.E. Freeman, and A.L. Kuck. 1991. Genetics of health traits in Holstein cattle. *J.*
 586 *Dairy Sci.* 74:1092–1100. doi:10.3168/jds.S0022-0302(91)78260-X.

587 Macrae, A.I., D.A. Whitaker, E. Burrough, A. Dowell, and J.M. Kelly. 2006. Use of metabolic
 588 profiles for the assessment of dietary adequacy in UK dairy herds. *Vet. Rec.* 159:655–661.
 589 doi:10.1136/vr.159.20.655.

590 Martinez, N., C.A. Risco, F.S. Lima, R.S. Bisinotto, L.F. Greco, E.S. Ribeiro, F. Maunsell, K.

591 Galvão, and J.E.P. Santos. 2012. Evaluation of peripartal calcium status, energetic profile,
592 and neutrophil function in dairy cows at low or high risk of developing uterine disease. *J.*
593 *Dairy Sci.* 95:7158–72. doi:10.3168/jds.2012-5812.

594 Masoero, F., M. Moschini, and A.M. Pulimeno. 2003. Serum calcium and magnesium level in
595 dairy cows at calving. *ITAL.J.ANIM.SCI.* 2 (Suppl. :172–174.

596 Melendez, P., G.A. Donovan, C.A. Risco, R. Littell, and J.P. Goff. 2003. Effect of calcium-
597 energy supplements on calving-related disorders, fertility and milk yield during the
598 transition period in cows fed anionic diets. *Theriogenology.* 60:843–854.
599 doi:10.1016/S0093-691X(03)00103-1.

600 Melendez, P., and C.A. Risco. 2005. Management of transition cows to optimize reproductive
601 efficiency in dairy herds. *Vet. Clin. North Am. - Food Anim. Pract.* 21:485–501.
602 doi:10.1016/j.cvfa.2005.02.008.

603 Mulligan, F.J., and M.L. Doherty. 2008. Production diseases of the transition cow. *Vet. J.* 176:3–
604 9. doi:10.1016/j.tvjl.2007.12.018.

605 Mulligan, F.J., L. O’Grady, D.A. Rice, and M.L. Doherty. 2006. A herd health approach to dairy
606 cow nutrition and production diseases of the transition cow. *Anim. Reprod. Sci.* 96:331–353.
607 doi:10.1016/j.anireprosci.2006.08.011.

608 NRC. 2001. Nutrient Requirements of Dairy Cattle Seventh Revised Edition , 2001. 1-333 pp.

609 Oetzel, G.R. 2011. Diseases of Dairy Animals | Non-Infectious Diseases: Milk Fever. *In*
610 *Encyclopedia of Dairy Sciences (Second Edition).* J.W. Fuquay, editor. Academic Press,
611 San Diego. 239–245.

612 Oikonomou, G., G. Arsenos, G.E. Valergakis, A. Tsiaras, D. Zygoyiannis, and G. Banos. 2008a.
613 Genetic relationship of body energy and blood metabolites with reproduction in Holstein

614 cows. *J. Dairy Sci.* 91:4323–4332. doi:10.3168/jds.2008-1018.

615 Oikonomou, G., G.E. Valergakis, G. Arsenos, N. Roubies, and G. Banos. 2008b. Genetic profile
616 of body energy and blood metabolic traits across lactation in primiparous Holstein cows. *J.*
617 *Dairy Sci.* 91:2814–2822. doi:10.3168/jds.2007-0965.

618 Peek, S.F., and T.J. Divers. 2008. Chapter 14 - Metabolic Diseases. *In* Rebhun's Diseases of
619 Dairy Cattle (Second Edition). T.J.D.F. Peek, editor. W.B. Saunders, Saint Louis. 590–605.

620 Pérez-Cabal, M.A., G. de los Campos, A.I. Vazquez, D. Gianola, G.J.M. Rosa, K.A. Weigel, and
621 R. Alenda. 2009. Genetic evaluation of susceptibility to clinical mastitis in Spanish Holstein
622 cows. *J. Dairy Sci.* 92:3472–80. doi:10.3168/jds.2008-1978.

623 Perkin Elmer. 1996. Perkin Elmer AAnalyst 100.

624 Pryce, J.E., R.F. Veerkamp, R. Thompson, W.G. Hill, and G. Simm. 1997. Genetic aspects of
625 common health disorders and measures of fertility in Holstein Friesian dairy cattle. *Anim.*
626 *Sci.* 65:353–360. doi:10.1017/S1357729800008559.

627 Reinhardt, T.A., J.D. Lippolis, B.J. McCluskey, J.P. Goff, and R.L. Horst. 2011. Prevalence of
628 subclinical hypocalcemia in dairy herds. *Vet. J.* 188:122–124.
629 doi:10.1016/j.tvjl.2010.03.025.

630 Risco, C.A. 2011. Management Considerations from Parturition to the End of the Voluntary
631 Waiting Period to Optimize Health and Reproductive Performance. *In* Dairy Production
632 Medicine. Blackwell Publishing Ltd. 1–6.

633 Rude, R.K. 1998. Magnesium deficiency: a cause of heterogeneous disease in humans. *J. Bone*
634 *Miner. Res.* 13:749–58. doi:10.1359/jbmr.1998.13.4.749.

635 Sattler, N., and G. Fecteau. 2014. Hypokalemia Syndrome in Cattle. *Vet. Clin. North Am. - Food*
636 *Anim. Pract.* 30:351–357. doi:10.1016/j.cvfa.2014.04.004.

637 Sattler, N., G. Fecteau, C. Girard, and Y. Couture. 1998. Description of 14 cases of bovine
 638 hypokalaemia syndrome. *Vet. Rec.* 143:503–507. doi:10.1136/vr.143.18.503.

639 Schonewille, J.T., H. Everts, S. Jittakhot, and A.C. Beynen. 2008. Quantitative prediction of
 640 magnesium absorption in dairy cows. *J. Dairy Sci.* 91:271–278. doi:10.3168/jds.2007-0304.

641 Sheldon, I.M., G.S. Lewis, S. LeBlanc, and R.O. Gilbert. 2006. Defining postpartum uterine
 642 disease in cattle. *Theriogenology*. 65:1516–1530.
 643 doi:10.1016/j.theriogenology.2005.08.021.

644 Staufenbiel, R. 2002. Neue Aspekte zum klinischen Bild und zur Therapie der Gebärpause des
 645 Rindes. *Vet Med Rev.* 26:12 V6.

646 Thilsing-Hansen, T., R.J. Jørgensen, and S. Østergaard. 2002. Milk fever control principles: a
 647 review. *Acta Vet. Scand.* 43:1–19. doi:10.1186/1751-0147-43-1.

648 Tsiamadis, V., G. Banos, N. Panousis, M. Kritsepi-Konstantinou, G. Arsenos, and G.E.
 649 Valergakis. 2016. Genetic parameters of calcium, phosphorus, magnesium, and potassium
 650 serum concentrations during the first 8 days after calving in Holstein cows. *J. Dairy Sci.*
 651 99:5535–5544. doi:10.3168/jds.2015-10787.

652 Uribe, H.A., B.W. Kennedy, S.W. Martin, and D.F. Kelton. 1995. Genetic parameters for
 653 common health disorders of Holstein cows. *J. Dairy Sci.* 78:421–430.
 654 doi:10.3168/jds.S0022-0302(95)76651-6.

655 Vazquez, A.I., K.A. Weigel, D. Gianola, D.M. Bates, M.A. Perez-Cabal, G.J.M. Rosa, and Y.M.
 656 Chang. 2009. Poisson versus threshold models for genetic analysis of clinical mastitis in US
 657 Holsteins. *J. Dairy Sci.* 92:5239–47. doi:10.3168/jds.2009-2085.

658 Veerkamp, R., and S. Brotherstone. 2001. Analysis of censored survival data using random
 659 regression models. *Anim. Sci.1994:* 1–10.






660 Windig, J.J., M.P.L. Calus, B. Beerda, and R.F. Veerkamp. 2006. Genetic correlations between
661 milk production and health and fertility depending on herd environment. *J. Dairy Sci.*
662 89:1765–1775. doi:10.3168/jds.S0022-0302(06)72245-7.

663 Wolf, V. 2001. Einflüsse auf das Auftreten von Labmagenverlagerungen bei Deutschen Holstein
664 Kühen - Influences on the occurrence of abomasal displacements in German Holstein cows.
665 *Dtsch. Tierärztl. Wochenschr.* 108.

666 Zwald, N.R., K.A. Weigel, Y.M. Chang, R.D. Welper, and J.S. Clay. 2004. Genetic selection for
667 health traits using producer-recorded data. II. Genetic correlations, disease probabilities, and
668 relationships with existing traits. *J. Dairy Sci.* 87:4295–4302. doi:10.3168/jds.S0022-
669 0302(04)73574-2.


670

GENETIC PARAMETRS OF HEALTH DISORDERS



Tsiamadis Figure 1.

GENETIC PARAMETRS OF HEALTH DISORDERS



Tsiamadis Figure 2.